

University of Groningen

## Penicillin-binding protein folding is dependent on the PrsA peptidyl-prolyl cis-trans isomerase in *Bacillus subtilis*

Hyyrylainen, Hanne-Leena; Marciniak, Bogumila C.; Dahncke, Kathleen; Pietiainen, Milla; Courtin, Pascal; Vitikainen, Marika; Seppala, Raili; Otto, Andreas; Becher, Doerte; Chapot-Chartier, Marie-Pierre

*Published in:*  
Molecular Microbiology

*DOI:*  
[10.1111/j.1365-2958.2010.07188.x](https://doi.org/10.1111/j.1365-2958.2010.07188.x)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2010

[Link to publication in University of Groningen/UMCG research database](#)

### *Citation for published version (APA):*

Hyyrylainen, H-L., Marciniak, B. C., Dahncke, K., Pietiainen, M., Courtin, P., Vitikainen, M., Seppala, R., Otto, A., Becher, D., Chapot-Chartier, M-P., Kuipers, O. P., Kontinen, V. P., Hyyryläinen, H-L., & Pietiäinen, M. (2010). Penicillin-binding protein folding is dependent on the PrsA peptidyl-prolyl cis-trans isomerase in *Bacillus subtilis*. *Molecular Microbiology*, 77(1), 108-127. <https://doi.org/10.1111/j.1365-2958.2010.07188.x>

### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# GFP-MreC is independent of PrsA




Fig.S1

# MreC is independent of PrsA




Fig.S1

# GFP-PBP2a levels in cell membranes




Fig. S1

# GFP-PBP1 is independent of PrsA




Fig. S1

Fig. S2






Fig. S3




**Fig. S3.** Identification of the main PBPs by null mutations of the PBP genes and Bocillin-FL staining (see also (Popham and Setlow, 1996)). Cytoplasmic membranes were isolated and PBPs were visualized by staining with Bocillin-FL (see Experimental procedures) and separation in SDS-PAGE. The identification of PBP2b is based on its known migration next to PBP2a (Popham and Setlow, 1996).

Popham, D. L., and Setlow, P. (1996) Phenotypes of *Bacillus subtilis* mutants lacking multiple class a high-molecular-weight penicillin-binding proteins. *J Bacteriol* **178**: 2079-2085.

A




B





C




**Fig. S4.** Quantitative analysis of PBP2a misfolding in PrsA-depleted cells. *cssS* mutant strains were used to prevent proteolytic degradation of the misfolded PBP2a, so that it could be identified and quantitated by Bocillin-FL staining (active PBP2a) and immunoblotting (total PBP2a: active and inactive).

A. Membrane samples were isolated from PrsA-depleted and non-depleted cells of *B. subtilis* IH9016 (*pbpA-myc Pspac-prsA cssS::spec*) and a series of two-fold diluted samples were subjected to SDS-PAGE and immunoblotted with anti-c-Myc antibodies. The amounts of full-length PBP2a-Myc were quantitated by determining optical densities of the PBP2a-Myc bands in lanes loaded with 7.5 µg and 3.7 µg of total membrane protein. The same dilution series of the membranes were also stained with Bocillin-FL and fluorescence intensities in full-length PBP2a-Myc were determined to quantitate correctly-folded active PBP2a-Myc.



B. The same analysis was also performed with *B. subtilis* IH8266 (*Pspac-prsA cssS::spec*) and anti-PBP2a antibodies.

C. Ratios of correctly-folded active PBP2a-Myc or PBP2a as determined by Bocillin-FL staining to total PBP2a-Myc or PBP2a as determined by immunoblotting were calculated, respectively. The ratios (level of active PBP2a-Myc or PBP2a) of non-depleted cells were adjusted to 100. The proportion of misfolded PBP2a-Myc or PBP2a in PrsA-depleted cells as compared to non-depleted cells was 30-40%. Fig. 3E suggests that the level of active PBP2a-Myc is lower in the PrsA-depleted *cssS* wild-type strain than in the PrsA-depleted *cssS* mutant, most probably implying that if misfolded PBP2a is not degraded due to the absence of the quality control proteases, at least a portion of this can slowly fold to an active conformation in the absence of the folding assistance of PrsA.

Fig. S5



# Small-angle X-ray scattering (SAXS) characterization of PrsA



**Fig.S6.** The radius of gyration ( $R_g$ ) is 27-29 Å, which is considerably larger than expected for a monomer. The measured  $R_g$  corresponds more like a tetramer.

**Table S2.** Plasmids used in this study.

Plasmid	Characteristics	Reference
pKTH3384	Derivative of pMUTIN4; Pspac- <i>'prsA'</i> ; Em <sup>r</sup>	(Vitikainen <i>et al.</i> , 2001)
pKTH3805	Derivative of pMUTIN-cMyc; <i>'prsA-myc'</i> ; Em <sup>r</sup>	This study
pKTH3806	Derivative of pSG4902; Pxyl- <i>gfp-divIC'</i> ; Cm <sup>r</sup>	This study
pKTH3814	Derivative of pSG4902; Pxyl- <i>gfp-divIB'</i> ; Cm <sup>r</sup>	This study
pKTH3828	Derivative of pSG4902; <i>'pbpA-myc'</i> ; Cm <sup>r</sup>	This study
pKTH3831	Derivative of pSG4902; <i>'dacA-myc'</i> ; Cm <sup>r</sup>	This study
pMUTIN-cMyc	Integration vector; Em <sup>r</sup>	(Kaltwasser <i>et al.</i> , 2002)
pSG4902	Integration vector; Cm <sup>r</sup>	(Scheffers <i>et al.</i> , 2004)

Kaltwasser, M., Wiegert, T., and Schumann, W. (2002) Construction and application of epitope- and green fluorescent protein-tagging integration vectors for *Bacillus subtilis*. *Appl Environ Microbiol* **68**: 2624-2628.

Scheffers, D. J., Jones, L. J., and Errington, J. (2004) Several distinct localization patterns for penicillin-binding proteins in *Bacillus subtilis*. *Mol Microbiol* **51**: 749-764.

Vitikainen, M., Pummi, T., Airaksinen, U., Wu, H., Sarvas, M., and Kontinen, V. P. (2001) Quantitation of the capacity of the secretion apparatus and requirement for PrsA in growth and secretion of  $\alpha$ -amylase in *Bacillus subtilis*. *J Bacteriol* **183**: 1881-1890.